

## § 331.11

## 21 CFR Ch. I (4–1–06 Edition)

(b) This section does not apply to an antacid ingredient specifically added as a corrective to prevent a laxative or constipating effect.

[39 FR 19874, June 4, 1974, as amended at 61 FR 4822, Feb. 8, 1996]

### § 331.11 Listing of specific active ingredients.

(a) Aluminum-containing active ingredients:

- (1) Basic aluminum carbonate gel.
- (2) Aluminum hydroxide (or as aluminum hydroxide-hexitol stabilized polymer, aluminum hydroxide-magnesium carbonate codried gel, aluminum hydroxide-magnesium trisilicate codried gel, aluminum-hydroxide sucrose powder hydrated).
- (3) Dihydroxyaluminum aminoacetate and dihydroxyaluminum aminoacetic acid.
- (4) Aluminum phosphate gel when used as part of an antacid combination product and contributing at least 25 percent of the total acid neutralizing capacity; maximum daily dosage limit is 8 grams.
- (5) Dihydroxyaluminum sodium carbonate.

(b) Bicarbonate-containing active ingredients: Bicarbonate ion; maximum daily dosage limit 200 mEq. for persons up to 60 years old and 100 mEq. for persons 60 years or older.

(c) Bismuth-containing active ingredients:

- (1) Bismuth aluminate.
- (2) Bismuth carbonate.
- (3) Bismuth subcarbonate.
- (4) Bismuth subgallate.
- (5) Bismuth subnitrate.
- (d) Calcium-containing active ingredients: Calcium, as carbonate or phosphate; maximum daily dosage limit 160 mEq. calcium (e.g., 8 grams calcium carbonate).
- (e) Citrate-containing active ingredients: Citrate ion, as citric acid or salt; maximum daily dosage limit 8 grams.
- (f) Glycine (aminoacetic acid).
- (g) Magnesium-containing active ingredients:

- (1) Hydrate magnesium aluminate activated sulfate.
- (2) Magaldrate.
- (3) Magnesium aluminosilicates.
- (4) Magnesium carbonate.
- (5) Magnesium glycinate.

(6) Magnesium hydroxide.

(7) Magnesium oxide.

(8) Magnesium trisilicate.

(h) Milk solids, dried.

(i) Phosphate-containing active ingredients:

(1) Aluminum phosphate; maximum daily dosage limit 8 grams.

(2) Mono or dibasic calcium salt; maximum daily dosage limit 2 grams.

(3) Tricalcium phosphate; maximum daily dosage limit 24 grams.

(j) Potassium-containing active ingredients:

(1) Potassium bicarbonate (or carbonate when used as a component of an effervescent preparation); maximum daily dosage limit 200 mEq. of bicarbonate ion for persons up to 60 years old and 100 mEq. of bicarbonate ion for persons 60 years or older.

(2) Sodium potassium tartrate.

(k) Sodium-containing active ingredients:

(1) Sodium bicarbonate (or carbonate when used as a component of an effervescent preparation); maximum daily dosage limit 200 mEq. of sodium for persons up to 60 years old and 100 mEq. of sodium for persons 60 years or older, and 200 mEq. of bicarbonate ion for persons up to 60 years old and 100 mEq. of bicarbonate ion for persons 60 years or older. That part of the warning required by § 330.1(g), which states, "Keep this and all drugs out of the reach of children" is not required on a product which contains only sodium bicarbonate powder and which is intended primarily for other than drug uses.

(2) Sodium potassium tartrate.

(l) Silicates:

(1) Magnesium aluminosilicates.

(2) Magnesium trisilicate.

(m) Tartrate-containing active ingredients. Tartaric acid or its salts; maximum daily dosage limit 200 mEq. (15 grams) of tartrate.

[39 FR 19874, June 4, 1974, as amended at 51 FR 27763, Aug. 1, 1986; 55 FR 19859, May 11, 1990]

### § 331.15 Combination with nonantacid active ingredients.

(a) An antacid may contain any generally recognized as safe and effective